

---

# Enhanced drug delivery via antibody–drug conjugates and siRNA-linked nucleic-acid aptamers targeting EGFR in glioblastoma cells

Charlène D'ancona\*<sup>1</sup>, Elisabete Cruz Da Silva<sup>1</sup>, Léa Denechere<sup>1</sup>, Hélène Justiniano<sup>1</sup>, Valérie Calco<sup>2</sup>, Romain Vauchelles<sup>1,3</sup>, Dilara Sensoy<sup>1</sup>, Candice Dussouillez<sup>4</sup>, Cendrine Seguin<sup>4</sup>, Pascal Villa<sup>2</sup>, Antoine Kichler<sup>4</sup>, Sylvie Fournel<sup>4</sup>, Halina Anton<sup>1</sup>, Maxime Lehmann<sup>1,5</sup>, and Laurence Choulier<sup>1</sup>

<sup>1</sup>Laboratoire de Bioimagerie et Pathologies – université de Strasbourg, Centre National de la Recherche Scientifique – France

<sup>2</sup>Plate-forme de chimie biologique intégrative de Strasbourg – université de Strasbourg, Centre National de la Recherche Scientifique – France

<sup>3</sup>Immunologie, Immunopathologie et Chimie Thérapeutique – Institut de biologie moléculaire et cellulaire, université de Strasbourg, Centre National de la Recherche Scientifique – France

<sup>4</sup>Centre de Recherche en Biomédecine de Strasbourg – université de Strasbourg, Institut National de la Santé et de la Recherche Médicale – France

<sup>5</sup>Etablissement Français du Sang - Grand Est – CNRS – France

## Résumé

Active targeting in drug delivery is based on the binding of ligands to receptors present on the surface of targeted cells in order to promote the internalization of ligand conjugated drugs. The most well-known conjugates are antibody-drug conjugates (ADCs), which combine the specificity of monoclonal antibodies with the cytotoxic of chemotherapeutic drugs. In addition to antibodies, nucleic acid aptamers, known as single stranded DNA or RNA oligonucleotides referring to chemical antibodies with high affinity and selectivity for their target (*Zhou et al, 2016*), are promising to deliver conjugated drugs or therapeutics nucleic acids such as siRNA by active targeting delivery (*Mercier et al, 2017*).

In our team, we are interested in developing an aptamer-siRNA chimera called AsiC targeting EGFR (epidermal growth factor receptor), a receptor, internalized by endocytosis (*Ivaska et al, 2011*) and often overexpressed in glioblastoma cells, the most aggressive tumour of the central nervous system. Our AsiC consists of a targeting part: a 2' fluoro modified RNA aptamer E07 targeting EGFR (Cruz da Silva, Foppolo et al, 2022) and a therapeutic part: a siRNA for gene silencing. The therapeutic efficacy of such conjugates not only depends on target specificity but also on efficient internalization into tumor cells. However, so far, no therapeutic approach to enhance endocytosis of conjugates is available.

In recent studies, we showed that gefitinib, a tyrosine kinase inhibitor directed against the EGFR, induces a massive, non-physiological endocytosis of EGFR, known as gefitinib-mediated endocytosis (GME), in different glioblastoma cell lines (*Blandin et al, 2021 ; Cruz Da Silva et al, 2021*). We thus hypothesized that besides promoting endocytosis of EGFR,

---

\*Intervenant

gefitinib could also promote endocytosis of its ligands. In this study, we proved by quantitative fluorescence bioimaging, that gefitinib is indeed able to strengthen the endocytosis of fluorophore-conjugated EGFR-specific antibodies and aptamers. We also showed that the GME potentiates the toxicity of an ADC and the efficacy of an AsiC. Our results suggest the development of a new therapeutic combination with gefitinib, to potentiate the delivery of ADC, AsiC and likely other conjugates targeting EGFR in glioblastoma, while limiting side effects on non-targeted cells. Our results have been submitted for publication and are already available online as a preprint (<https://www.biorxiv.org/content/10.1101/2024.10.22.617611v1>).

Blandin, A.-F., Cruz Da Silva, E., Mercier, M.-C., Glushonkov, O., Didier, P., Dedieu, S., Schneider, C., Devy, J., Etienne-Selloum, N., Dontenwill, M., Choulier, L., Lehmann, M., 2021. Gefitinib induces EGFR and  $\alpha 5\beta 1$  integrin co-endocytosis in glioblastoma cells. *Cell. Mol. Life Sci.* 78, 2949–2962. <https://doi.org/10.1007/s00018-020-03686-6>

Cruz Da Silva, E., Choulier, L., Thevenard-Devy, J., Schneider, C., Carl, P., Rondé, P., Dedieu, S., Lehmann, M., 2021. Role of Endocytosis Proteins in Gefitinib-Mediated EGFR Internalisation in Glioma Cells. *Cells* 10, 3258. <https://doi.org/10.3390/cells10113258>

Cruz Da Silva, E.\*, Foppolo, S.\*, Lhermitte, B., Ingremeau, M., Justiniano, H., Klein, L., Chenard, M.-P., Vauchelles, R., Abdallah, B., Lehmann, M., Etienne-Selloum, N., Dontenwill, M., Choulier, L. (2022). Bioimaging nucleic acid aptamers of different specificities on human glioblastoma tissues highlights tumoral heterogeneity. *Pharmaceutics* 14(10), 1980. doi: 10.3390/pharmaceutics14101980 (\*co-first authors)

Ivaska J, Heino J. Cooperation between integrins and growth factor receptors in signaling and endocytosis. *Annu Rev Cell Dev Biol.* 2011;27:291-320. doi:10.1146/annurev-cellbio-092910-154017

Mercier, M.-C., Dontenwill, M., Choulier, L., 2017. Selection of Nucleic Acid Aptamers Targeting Tumor Cell-Surface Protein Biomarkers. *Cancers (Basel)* 9. <https://doi.org/10.3390/cancers9060069>

Zhou, G., Wilson, G., Hebbard, L., Duan, W., Liddle, C., George, J., Qiao, L., 2016. Aptamers: A promising chemical antibody for cancer therapy. *Oncotarget* 7, 13446–13463. <https://doi.org/10.18632/oncotarget.7178>

**Mots-Clés:** antibody, drug conjugates, bioimaging, epidermal growth factor, gefitinib, glioblastoma, endocytosis, nucleic acid aptamers